

Differential geometry of fibre bundles over foliated manifolds

G.SARDANASHVILY

Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia

E-mail: sard@grav.phys.msu.su

Abstract. Any leafwise connection on a fibre bundle over a foliated manifold is proved to come from a connection on this fibre bundle.

Mathematics Subject Classification (2000): 53C05, 53C12

Let $(Z, \mathcal{F})$ be a foliated manifold, where $\mathcal{F}$ denotes a regular foliation of Z . Let $Y \rightarrow Z$ be a fibre bundle over Z . A leafwise connection on $Y \rightarrow Z$ is defined as a partial connection along the leaves of a foliation $\mathcal{F}$. Obviously, each connection on $Y \rightarrow Z$ yields a leafwise connection. We show that any leafwise connection on $Y \rightarrow Z$ is of this type. This fact is important, e.g., for geometric quantization of symplectic foliations.

We start from the notion of a leafwise differential calculus on a foliated manifold. All manifolds throughout are smooth, real, finite-dimensional, Hausdorff, paracompact, and connected.

If $(Z, \mathcal{F})$ is a (regular) foliated manifold, there is an adapted coordinate atlas

$$\Psi_{\mathcal{F}} = \{(U; z^{\alpha}; z^A)\}, \quad \alpha = 1, \dots, \dim \mathcal{F}, \quad A = \dim \mathcal{F} + 1, \dots, \dim Z, \quad (1)$$

of Z such that, for every leaf F of a foliation $\mathcal{F}$, the connected components of $F \cap U$ are described by the equations $z^A = \text{const.}$, and transition functions of coordinates z^A are independent of the remaining ones [3].

The tangent spaces to leaves of a foliation $\mathcal{F}$ assemble into an involutive distribution $T\mathcal{F} \rightarrow Z$ on Z called the tangent bundle to $\mathcal{F}$. Its global sections constitute the real Lie algebra $T\mathcal{F}(Z)$ of vector fields on Z subordinate to $T\mathcal{F}$. This Lie algebra is a left $C^{\infty}(Z)$ -submodule of the derivation module of the $\mathbb{R}$ -ring $C^{\infty}(Z)$ of smooth real functions on Z . Let $S_{\mathcal{F}}(Z)$ be the kernel of $T\mathcal{F}(Z)$, acting on $C^{\infty}(Z)$. It consists of functions, constant on leaves of $\mathcal{F}$ (i.e., foliated functions in the terminology of [4]). Then $T\mathcal{F}(Z)$ is the Lie $S_{\mathcal{F}}(Z)$ -algebra of derivations of $C^{\infty}(Z)$, seen as a $S_{\mathcal{F}}(Z)$ -ring. Let us consider the Chevalley–Eilenberg differential calculus over the $S_{\mathcal{F}}(Z)$ -ring $C^{\infty}(Z)$. It is defined as a cochain subcomplex

$$0 \rightarrow S_{\mathcal{F}}(Z) \longrightarrow C^{\infty}(Z) \xrightarrow{\tilde{d}} \mathfrak{F}^1(Z) \xrightarrow{\tilde{d}} \dots \xrightarrow{\tilde{d}} \mathfrak{F}^{\dim \mathcal{F}}(Z) \rightarrow 0 \quad (2)$$

of the Chevalley–Eilenberg complex of the Lie $S_{\mathcal{F}}(Z)$ -algebra with coefficients in the ring $C^\infty(Z)$, where $\mathfrak{F}^r(Z)$, $r > 0$, is a $C^\infty(Z)$ -module of $C^\infty(Z)$ -multilinear skew-symmetric maps

$$\phi : \times^r T\mathcal{F}(Z) \rightarrow C^\infty(Z)$$

[2]. These maps are precisely global sections of the exterior product $\bigwedge^r T\mathcal{F}^*$ of the dual $T\mathcal{F}^* \rightarrow Z$ of $T\mathcal{F} \rightarrow Z$. They are called the leafwise forms on a foliated manifold $(Z, \mathcal{F})$ (one should distinguish them from foliated forms in [4] which are exterior forms on Z constant on leaves of a foliation). With respect to the adapted coordinates $(z^\alpha; z^A)$ (1), the leafwise forms read

$$\phi = \frac{1}{r!} \phi_{\alpha_1 \dots \alpha_r} \tilde{d}z^{\alpha_1} \wedge \dots \wedge \tilde{d}z^{\alpha_r}, \quad (3)$$

where $\{\tilde{d}z^\alpha\}$ are the duals of the holonomic fibre bases $\{\partial_\alpha\}$ for $T\mathcal{F}$. The Chevalley–Eilenberg coboundary operator on leafwise forms is given by the expression

$$\tilde{d}\phi = \tilde{d}z^\mu \wedge \partial_\mu \phi = \frac{1}{r!} \partial_\mu \phi_{\alpha_1 \dots \alpha_r} \tilde{d}z^\mu \wedge \tilde{d}z^{\alpha_1} \wedge \dots \wedge \tilde{d}z^{\alpha_r}, \quad \mu, \alpha \leq \dim \mathcal{F}, \quad (4)$$

(it is precisely the operator d_f in [4]). It is called the leafwise differential, and $(\mathfrak{F}^*(Z), \tilde{d})$ is the leafwise differential calculus [1].

Let $V\mathcal{F} = TZ/T\mathcal{F} \rightarrow Z$ be the normal bundle to a foliation $\mathcal{F}$ and $V\mathcal{F}^*$ the dual of $V\mathcal{F}$. There are the exact sequences of vector bundles

$$0 \rightarrow T\mathcal{F} \xrightarrow[Z]{i} TZ \xrightarrow[Z]{j} V\mathcal{F} \rightarrow 0, \quad (5)$$

$$0 \rightarrow V\mathcal{F}^* \xrightarrow[Z]{j^*} T^*Z \xrightarrow[Z]{i^*} T\mathcal{F}^* \rightarrow 0. \quad (6)$$

The epimorphism i^* in (6) provides an order-preserving epimorphism of the graded differential algebra $(\mathcal{O}^*(Z), d)$ of exterior forms on Z to the algebra $(\mathcal{F}^*(Z), \tilde{d})$ of leafwise forms such that $i^* \circ d = \tilde{d} \circ i^*$. With respect to the adapted coordinates (1), this epimorphism reads

$$i^* : dz^\alpha \mapsto \tilde{d}z^\alpha, \quad i^* : dz^A \rightarrow 0.$$

Turn now to the notion of a leafwise connection on a fibre bundle

$$\pi : Y \rightarrow Z \quad (7)$$

over a foliated manifold $(Z, \mathcal{F})$. The inverse images $\pi^{-1}(F)$ of leaves F of the foliation $\mathcal{F}$ of Z provides a (regular) foliation $Y_{\mathcal{F}}$ of Y . One can always choose an adapted coordinate

atlas $\{(U; z^\alpha; z^A)\}$ (1) of a foliated manifold $(Z, \mathcal{F})$ such that U are trivialization domains of a fibre bundle $Y \rightarrow Z$. Let $(z^\alpha; z^A; y^i)$ be the corresponding bundle coordinates on $Y \rightarrow Z$. They are also adapted coordinates on the foliated manifold $(Y, Y_{\mathcal{F}})$.

Let us note that, in the terminology of [4], a fibre bundle $Y \rightarrow Z$ (7) is called foliated if its transition functions are constant along leaves of $\mathcal{F}$.

Given the tangent bundle $TY_{\mathcal{F}} \rightarrow Y$ to the foliation $Y_{\mathcal{F}}$ of Y , we have the exact sequence of vector bundles

$$0 \rightarrow VY \xrightarrow{Y} TY_{\mathcal{F}} \xrightarrow{Y} Y \times_Z T\mathcal{F} \rightarrow 0, \quad (8)$$

where VY is the vertical tangent bundle of $Y \rightarrow Z$. Its splitting exists, and is called a leafwise connection on a fibre bundle $Y \rightarrow Z$. It is represented by a $TY_{\mathcal{F}}$ -valued leafwise one-form

$$A_{\mathcal{F}} = \tilde{d}z^\alpha \otimes (\partial_\alpha + A_\alpha^i \partial_i). \quad (9)$$

Let $J_{\mathcal{F}}^1 Y$ be the subbundle of the tensor product $T\mathcal{F}^* \otimes_Y TY_{\mathcal{F}} \rightarrow Y$ whose image under the surjection

$$T\mathcal{F}^* \otimes_Y TY_{\mathcal{F}} \xrightarrow{Y} Y \times_Z (T\mathcal{F}^* \otimes T\mathcal{F})$$

coincides with the canonical section $\theta_{\mathcal{F}}(Y) = \tilde{d}z^\alpha \otimes \partial_\alpha$ of the pull-back

$$Y \times_Z (T\mathcal{F}^* \otimes T\mathcal{F}) \rightarrow Y.$$

One can think of $J_{\mathcal{F}}^1 Y$ as being a leafwise jet manifold of a fibre bundle $Y \rightarrow Z$. With respect to the adapted bundle coordinates $(z^\alpha; z^A; y^i)$, its elements read

$$\tilde{d}z^\alpha \otimes (\partial_\alpha + y_\alpha^i \partial_i).$$

In particular, every section s of a fibre bundle $Y \rightarrow Z$ gives rise to the section

$$J_{\mathcal{F}}^1 s = \tilde{d}z^\alpha \otimes (\partial_\alpha + \partial_\alpha s^i \partial_i)$$

of the leafwise jet bundle $J_{\mathcal{F}}^1 Y \rightarrow Z$.

Similar to the case of a connection, one can show that every leafwise connection on a fibre bundle $Y \rightarrow Z$ is a global section of the affine leafwise jet bundle

$$\pi_0^1 : J_{\mathcal{F}}^1 Y \rightarrow Y,$$

and *vice versa*. It follows that leafwise connections constitute an affine space modelled on the vector space of sections of the vector bundle

$$T\mathcal{F}^* \otimes_Y VY \rightarrow Y. \quad (10)$$

In particular, we have the leafwise covariant differential

$$\nabla_{\mathcal{F}} = \text{Id} - A_{\mathcal{F}} \circ \pi_0^1 : J_{\mathcal{F}}^1 Y \rightarrow T\mathcal{F}^* \otimes_Y VY,$$

$$\nabla_{\mathcal{F}} s = (\tilde{d}s^i - (A_{\alpha}^i \circ s) \tilde{d}z^{\alpha}) \otimes \partial_i.$$

Since the exact sequence (8) is a subsequence of the exact sequence

$$0 \rightarrow VY \xrightarrow{Y} TY \xrightarrow{Y} Y \times_Z TZ \rightarrow 0,$$

every connection

$$\Gamma = dz^A \otimes (\partial_A + \Gamma_A^i \partial_i) + dz^{\alpha} \otimes (\partial_{\alpha} + \Gamma_{\alpha}^i \partial_i)$$

on a fibre bundle $Y \rightarrow Z$ yields a leafwise connection

$$\Gamma_{\mathcal{F}} = \tilde{d}z^{\alpha} \otimes (\partial_{\alpha} + \Gamma_{\alpha}^i \partial_i). \quad (11)$$

Conversely, let $A_{\mathcal{F}}$ (8) be a leafwise connection on a fibre bundle $Y \rightarrow Z$ and $\Gamma_{\mathcal{F}}$ (11) a leafwise connection which comes from a connection on $Y \rightarrow Z$. Their affine difference over Y is a section

$$Q = A_{\mathcal{F}} - \Gamma_{\mathcal{F}} = \tilde{d}z^{\alpha} \otimes (A_{\alpha}^i - \Gamma_{\alpha}^i) \partial_i$$

of the vector bundle (10). Given some splitting

$$B : \tilde{d}z^{\alpha} \mapsto dz^{\alpha} - B_A^{\alpha} dz^A \quad (12)$$

of the exact sequence (6), the composition

$$B \circ Q : Y \rightarrow T^*Z \otimes_Y VY,$$

$$B \circ Q = (dz^{\alpha} - B_A^{\alpha} dz^A) \otimes (A_{\alpha}^i - \Gamma_{\alpha}^i) \partial_i,$$

is a soldering form on a fibre bundle Y . Then

$$\Gamma + B \circ Q = dz^A \otimes (\partial_A + \Gamma_A^i \partial_i - B_A^{\alpha} (A_{\alpha}^i - \Gamma_{\alpha}^i) \partial_i) + dz^{\alpha} \otimes (\partial_{\alpha} + A_{\alpha}^i \partial_i)$$

is a desired connection on a fibre bundle $Y \rightarrow Z$ which yields the leafwise connection $A_{\mathcal{F}}$ (8).

Of course, such a connection is not unique, but depends on the choice of a trivialization B (12) of the exact sequence (6).

References

- [1] Hector, G., Macías, E. and Saralegi, M.: Lemme de Moser feuilleté et classification des variétés de Poisson régulières, *Publ. Mat.* **33** (1989), 423.
- [2] Mangiarotti, L. and Sardanashvily, G.: *Connections in Classical and Quantum Field Theory*, World Scientific, Singapore, 2000.
- [3] Reinhart, B.: *Differential Geometry and Foliations*, Springer-Verlag, Berlin, 1983.
- [4] Vaisman, I.: *Cohomology and Differential Forms*, Marcel Dekker, Inc., New York, 1973.